

## Anxiety and Depression among Nursing Staff at King Fahad Medical City, Kingdom of Saudi Arabia

Mostafa A F. Abbas<sup>1,2</sup>; Lamiaa Z. Abu Zaid<sup>1,2</sup>; Mona Hussaein<sup>3</sup>; Khaled H. Bakheet<sup>1</sup> and Nasser A. AlHamdan<sup>1</sup>

<sup>1</sup> Faculty of Medicine; King Saud Bin Abdulaziz University for Health Science (KSAU-HS), King Fahad Medical City (KFMC), Riyadh, Kingdom of Saudi Arabia.

<sup>2</sup> Faculty of Medicine, Suez Canal University, Ismailia, Egypt.

<sup>3</sup> Nursing Administration, KFMC, Riyadh, Kingdom of Saudi Arabia.

### Abstract:

**Background:** Nursing staff are exposed to numerous stresses that affect their psychological status. This study aimed to survey nursing staff at KFMC, for anxiety and depression symptoms using HAD scale. **Methods:** Nursing staff at KFMC were received self-administered questionnaires containing socio-demographic data, work related data and the English and Arabic versions of Hospital Anxiety and Depression scale (HADS). **Results:** For anxiety 53% of the study subjects were normal (scores 0-7) and 27% were classified as cause of concern (scores 8-10) while the probable clinical cases (scores  $\geq 11$ ) represented 20%. For depression, 75% were normal (scores 0-7) and 15% were classified as cause of concern (scores 8-10) while the probable clinical cases (scores  $\geq 11$ ) represented 10%. The highest prevalence rate of probable clinical cases of anxiety (23%) was reported among the age group 20 to less than 30 years. While the highest prevalence rate of probable clinical cases of depression (10.9%) was reported among the age group 30 to less than 40 years. Nurses from the Middle East region have significantly higher anxiety and depression symptoms prevalence rates than the other nationalities. Also, nurses who are not practicing physical activity have significantly higher anxiety and depression symptoms prevalence rates than those who are practicing physical activity. For smoking, there was a significant association between both anxiety and depression and the current smoking status where smokers have higher anxiety and depression symptoms prevalence rates. **Conclusion:** Middle Eastern nurses, Divorced/widowed nurses, lack of physical exercise and smoking were risk factors for anxiety/depression symptoms among nursing staff.

**Keywords :** nursing staff; HAD scale; anxiety; depression

### Introduction:

Depression and anxiety are common conditions with prevalence ranging between ten to twenty percent in the general population for any twelve-month period.<sup>1</sup> Depression in the working age population is estimated to cost \$12 billion annually in medical care and approximately \$44 billion annually in lost productivity.<sup>2</sup> Mental health affects physical health, job performance and healthcare utilization. Stress, depression and anxiety disorders contribute to absenteeism and lack of confidence<sup>3</sup>. Hospital anxiety and depression scale is a self assessment tool has been developed to be a reliable instrument for detecting states of depression and anxiety in the setting of hospital, medical or outpatient clinic.<sup>4-6</sup> There is a strong international experience with use of HAD scale of more than 200 published studies from most medical settings worldwide. The HADS gives clinically meaningful results as a psychological screening tool, in clinical groups comparisons and in correlation studies with several aspects of disease and quality of life.<sup>7-8</sup> Several studies provide evidence for the association between work stress and mild psychiatric morbidity among emergency nurses. The most commonly reported manifestations were sleep disturbances, anxious mood, and depressed mood.<sup>9</sup> A descriptive, correlation study was performed in Brazil<sup>10</sup> with the objective to evaluate anxiety and depression among nursing professionals working in surgical units. Participants were 211 nurses and data collection was performed using a questionnaire for the socio-demographic characterization and the Hospital Anxiety and Depression Scale. Results showed that the workers average score. Another study conducted in Greece for comparison of anxiety and depression for 76 Greek nursing personnel and 66 doctors using validated translations of the Spielberger State-Trait Anxiety Inventory and the Beck Depression Inventory. Male nurses had the lowest scores on trait anxiety, while female doctors had the highest scores, followed by female nurses. Age and depression scores were positively correlated only for the female nurses. The findings suggest that interventions for diagnosis and stress management in the health-care workplace should be equally

targeted for nursing and medical personnel.<sup>11</sup> Anxiety and depression are the most common psychological disorders that affect worker population. It can affect the productivity of the institution. Addressing them and its associated risk factors will help reduce these disorders among working population that will improve productivity, reduce absence and loss of resources on treatment, investigation and rehabilitation. This study aimed to determine the prevalence rate of and associated factors for anxiety and depression symptoms among nursing staff.

#### Materials and Methods:

**Study design:** this is a cross sectional study, conducted at KFMC, a tertiary care health facility in Riyadh, Saudi Arabia.



**Population and sample:** the study population comprised nursing staff from all hospitals and centers inside KFMC, 1300 questionnaires and HAD scale were distributed; participation was optional and 715 (55%) nurses had been participated in this study. Participants' consent and ethical approval were obtained; IRB number 11-086.

**Data collection instruments:** We used a questionnaire covering socio-demographic data. We also used the HAD scale, both English and Arabic versions, the Arabic one was translated and validated by Dwedat.<sup>12</sup> The scale contains 14 multiple-choice questions, with two sub-scales: anxiety (HADS-A) and depression (HADS-D), with seven items in each domain. Scores for each item range from zero to three, and the global score for each subscale range from zero to 21. To interpret the scores of the two subscales, it is considered that, the higher the score, the greater the chance that the person will develop an anxiety and/or depression disorder.<sup>4</sup> The cut-off points for normal score is equal to or less than 7, 8-10 will be classified as cause of concern – monitor for change, scores more than or equal to 11, will be tagged as probable clinical case. **Statistical analysis:** Collected data were processed using SPSS software, frequency distribution tables and pie-charts were produced. Comparison of continuous variables among different groups was done using students t test. Cross tables for different categories of categorical variables were done. Chi square test was used for test of difference among different groups for categorical data. Multivariate logistic regression analysis was used to check for the effect of multiple variables on occurrence of anxiety and depression.

**Table (1): Socio-demographic and Working Life Characteristics of the Study Subjects (n=715)**

Variable	Frequency	Percent
<i>Age in years (mean ± SD)</i>	35.2 ± 8.2	
<b>Gender:</b>		
Male	83	11.6%
Female	632	88.4%
<b>Marital status:</b>		
Married	492	68.8%
Single	198	27.6%
Widow	13	1.8%
Divorced	12	1.6%
<b>Nationality:</b>		
South Asia	527	73.7%

Middle East	111	15.5%
African	48	6.7%
Western	29	4.0%
<b>Physical Activity:</b>		
Yes	465	65.0%
No	250	35.0%
<b>Smoking:</b>		
No	655	91.6%
Yes	60	8.4%
<b>Body Mass Index:</b> (mean ± SD)		
24.5 ± 4.4		
<b>Work duration/year at KFMC:</b> (mean ± SD)		
4.3 ± 2.3		
<b>Working hours/day:</b> (mean ± SD)		
11.5 ± 0.8		
<b>Night shift:</b>		
Yes	508	71.0%
No	207	29.0%
<b>Overtime work:</b>		
Yes	56	7.8%
No	659	92.2%



**Table (2): Association between Socio-demographic Characteristics and Presence of Anxiety (Probable Clinical Case, Scores ≥ 11) among Nursing Staff at KFMC, Saudi Arabia, 2011**

Socio-demographic Characteristics	No anxiety (scores < 11) n= 574 No. (%)	Probable clinical case of anxiety (scores ≥11) n= 141 No. (%)	P-value
<b>Age (yr):</b>			
20-	161 (77.0%)	48 (23.0%)	0.28
30-	258 (80.0%)	62 (20.0%)	
40-	155 (83.3%)	31 (16.7%)	

**Table (3): Association between Socio-demographic Characteristics and Presence of Depression (Probable Clinical Case, Scores  $\geq 11$ ) among Nursing Staff at KFMC, Saudi Arabia, 2011**

<i>Socio-demographic Characteristics</i>	No depression (scores < 11) n= 646 No. (%)		Probable clinical case of depression (scores $\geq 11$ ) n= 69 No. (%)		<i>P-value</i>
	<i>Age (yr):</i>				
20-	185 (89.8%)		21 (10.2%)		0.34
30-	288 (89.1%)		35 (10.9%)		
40-	173 (91.0%)		13 (9.0%)		
<i>Gender:</i>					0.0000
Male	61 (73.5%)		22 (26.5%)		
Female	585 (92.6%)		47 (7.4%)		
<i>Marital status:</i>					0.016
Married	448 (91.2%)		44 (8.8%)		
Single	180 (90.6%)		18 (9.4%)		
Widow	9 (81.8%)		4 (18.2%)		
Divorced	9 (90.0%)		3 (10%)		
<i>Nationality:</i>					0.0000
South Asia	505 (95.8%)		22 (4.2%)		
Middle East	76 (68.4%)		35 (31.6%)		
African	38 (79.2%)		10 (20.8%)		
Western	27 (93.0%)		2 (7.0%)		
<i>Physical Activity:</i>					0.036
Yes	428 (92.0%)		37 (8.0%)		
No	218 (87.2%)		32 (12.8%)		
<i>Smoking</i>					

**Table (5): Logistic Regression Analysis for Anxiety as a Dependent Variable**

Variable	Beta	Standard Error	P- value	Odds Ratio	95% C.I. for OR
<i>Smoking</i>					
Yes	0.66	1.12	0.54	1.94	0.24-15.87
No	-1.04	0.33	0.00	0.36	0.19- 0.67
<i>Physical Activity</i>					
Yes	-0.63	0.56	0.26	0.53	0.18- 1.59
No	0.43	0.22	0.04	1.53	1.00- 2.34
<i>Constant</i>	-0.66	0.31	0.03	0.52	

**Table (6): Logistic Regression Analysis for Depression as a Dependent Variable**

Variable	Beta	Standard Error	P- value	Odds Ratio	95% C.I. for OR
<i>Smoking</i>					
Yes	-20.11	170	0.99	0.00	0
No	-1.12	0.43	0.01	0.33	0.14-0.76
<i>Work duration/ year at KPMC</i>	0.14	0.07	0.04	1.14	1.00- 1.31
<i>Constant</i>	2.89	2.18	0.18	18.12	

**Discussion:**

Nursing staff are exposed to a variety of work related stress that affects their productivity and their psychological health. This study showed that 47% of nursing staff had anxiety symptoms (HAD Score  $\geq 8$ ). Of those, 20% were probable clinical cases (HAD Score  $\geq 11$ ). These figures are higher than the prevalence reported by Schmidt D. et al., 2011,<sup>10</sup> in a study conducted in Brazil and found that the prevalence of anxiety (HAD Score  $\geq 8$ ) was 31.3% among nursing professionals working at surgical units. In another study conducted in Greece<sup>9</sup> and surveyed 213 nursing staff to investigate the presence of anxiety and stress symptoms among emergency nursing personnel, using the Hamilton anxiety scale, the authors reported that 10.7% had very severe anxious mood. The differences in reported prevalence might be explained by the use of a different survey tool than ours, also we had a much larger sample size and we have to put into consideration the diversity of the nationalities in our study sample of nursing staff at KFMC. As for depression, in the current study 25% of nursing staff had symptoms of depression (HAD score  $\geq 8$ ), of those, only 10% represented probable clinical case (HAD score  $\geq 11$ ), which are comparable to the study conducted by Schmidt D; 2011<sup>10</sup> who reported that the prevalence of depression symptoms among nursing professionals were 24.2%. While in the Stathopoulou study in Greece<sup>9</sup> nursing staff had a high prevalence (23.9%) of severe depressive mood, which is even much higher figure than our study, this difference might be explained by use of different survey tool. In the present study, we couldn't find statistically significant association between both overtime work and night shift and the manifestations of anxiety (HAD Score  $\geq 11$ ) or depression as shown in Table 4; which is in contrast to the study of Boya F et al., 2008<sup>13</sup>, the authors reported that anxiety is significantly correlated with both night work and overtime work. In our study, we found that nurses who are practicing physical exercise are less likely to have anxiety symptoms compared to those who are not practicing physical exercise, using chi square test (Table 2,  $p=0.034$ ). Also interestingly, in our study we found that nursing staff who are smokers are more likely to have anxiety and depression symptoms (tables 2 and 3). In multivariate analysis we found significant association between smoking and depression symptoms using HAD scale (Table 6). Mykletun A et al., 2007<sup>14</sup>, examined association of smoking with depression and anxiety, he screened 60814 in a population-based health survey using the HAD scale and reported that the association with smoking was strongest in co-morbid anxiety depression, followed by anxiety, and only marginal in depression. Findings of Mykletun A et al., 2007<sup>14</sup> is in agreement with our findings inspite of our much smaller sample size compared to his, and it was a single center study.

**Conclusion and Recommendations:**

Middle Eastern nurses, Divorced/widowed nurses, lack of physical exercise and smoking were risk factors for anxiety/depression symptoms among nursing staff. Major health education and health promotion programs are required to foster exercise and no smoking culture among nursing staff. Annual HAD scale might be helpful in identifying nursing staff who are considered as probable clinical case of anxiety and/or depression for support program from nursing administration.

**Biography of each author:**

- Mostafa A F. Abbas, MBBS, MD, PhD, MSc. Medical Education. Professor of Occupational Medicine, Faculty of Medicine, Suez Canal University, Egypt. E-mail: [moustafafouad@yahoo.com](mailto:moustafafouad@yahoo.com)
- Lamiaa Z. Abu Zaid, MBBS, MD, PhD. Assistant Professor of Community Medicine, faculty of Medicine, KFMC, KSAU-HS, Kingdom of Saudi Arabia. E-mail: [labuzaid@kfmc.med.sa](mailto:labuzaid@kfmc.med.sa)
- Nasser A. AlHamdan, MBBS, FFCM. Consultant and Associate Professor of Family and Community medicine, faculty of Medicine, KFMC, , KSAU-HS, Kingdom of Saudi Arabia. E-mail: [nalhamdan@kfmc.med.sa](mailto:nalhamdan@kfmc.med.sa)
- Mona Hussaein, Bsc, MSc., Nursing Administration, King Fahad Medical City, Kingdom of Saudi Arabia. E-mail: [mhusssein@kfmc.med.sa](mailto:mhusssein@kfmc.med.sa)
- Khaled H. Bakheet, 6<sup>th</sup> year medical student, faculty of Medicine, KFMC, KSAU-HS, Kingdom of Saudi Arabia. E-mail: [khalid\\_50\\_1@hotmail.com](mailto:khalid_50_1@hotmail.com)

**References:**

1. Center C, Davis M, Detre T, Ford D, et al. Confronting depression and suicide in physicians: A consensus statement. *JAMA*. 2003;289:3161-3166.
2. Kessler RC, McGonagle KA, Zhao S et al. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in US: Results from the National Comorbidity Survey. *Arch General Psychiatry*. 1994;51:8-19.
3. Williams S, Dale J, Glucksman E, Wellesley A. Senior house officers' work related stressors, psychological distress, and confidence in performing clinical tasks in accident and emergency: a questionnaire study. *BMJ*. 1997;713:718.
4. Zigmond AS and Snaith RP: The hospital anxiety and depression scale. *Acta Psychiatr Scand*. 1983 Jun; 67(6): 361-70
5. Zigmond AS and Snaith RP: the hospital anxiety and depression scale, *BMJ* 1986, (292) p 344.
6. Zigmond AS and Snaith RP: anxiety and depression in general medical settings. *BMJ* 1988, (297), p 1544.
7. Herrmann C.: International experiences with the hospital anxiety and depression scale- a review of validation data and clinical results. *J Psychosom Res*. 1997, Jan; 42(1):17-41
8. Snaith P: the hospital anxiety and depression scale. *Health and quality of life outcomes (OpenAccess)*, 2003 p1-29.
9. Stathopoulou H, Karanikola MN, Panagiotopoulou F, Papatheanassoglou ED: Anxiety Levels and Related Symptoms in Emergency Nursing Personnel in Greece. *J Emerg Nurs*. 2010 Apr 21. [Epub ahead of print]
- 10 Schmidt D; Dantas R; and Marziale M: Anxiety and depression among nursing professionals who work in surgical units. *Rev Esc Enferm USP* 2011; 45(2):475-81.
11. Athanassios Tselebis, Georgios Gournas, Georgia Tzitzanidou, Aspasia Panagiotou And Ioannis Ilias : Anxiety And Depression In Greek Nursing And Medical Personnel. *Psychological Reports (2006): Volume 99, Issue , pp. 93-96.*
- 12 Dwedat A.: The hospital anxiety and depression scale. Arabic translation and validation of the scale. Alexandria university, Faculty of Literature. Psychology dept.
13. Boya F; Demiral Y; Ergor A; Akvardary Y and Witte H: effects of perceived job insecurity on perceived anxiety and depression in nurses . *Indust Health* 2008, 46, 613-619.
14. Mykletun A; Overland S; Aaro L; Liabo H and Stewart R.: smoking in relation to anxiety and depression: evidence from a large population survey. *The HUNT study. European Psychiat*. 2008 (23)77-84.